misestudy.com smart learning



IIT-JEE · NEET · CBSE eBOOKS

CLASS 11&12th



CLASS 12th
Electrodynamics



01. Magnetic Field

Magnetic Field

The magnetic field is a space around a conductor carrying current or the space around a magnet in which its magnetic effect can be felt.

Moving charge is a source of both electric field as well as a magnetic field. Magnetic field denoted by \overrightarrow{B} is a vector.

To define the magnetic field \overrightarrow{B} , we deduce an expression for the force on a moving charge in a magnetic field.


$$F \propto q v \sin \theta B$$
 or $F = k q v B \sin \theta$

Where k is a constant

$$|\vec{F}| = q |\vec{v} \times \vec{B}|$$
 or $\vec{F} = q (\vec{v} \times \vec{B})$


Direction of \overrightarrow{F}

Given by the Right-Handed-Screw rule or Right-Hand Rule.



02. Biot-Savart's Law

This law deals with the magnetic field induction at a point due to a small current element.



In SI Units,

$$d\mathbf{B} = \frac{\mu_0}{4\pi} \times \frac{Idl \sin \theta}{r^2}$$

In vector form,

$$\overrightarrow{dB} = \frac{\mu_0}{4\pi} \frac{I(\overrightarrow{dl} \times \overrightarrow{r})}{r^3}$$

Direction of $d\vec{B}$

Right handed screw rule or Right Hand Rule.


Biot Savart's law in terms of charge (q) and its velocity (v) is.

$$\vec{dB} = \frac{\mu_0}{4\pi} \frac{q(\vec{v} \times \vec{r})}{r^3}$$

Important features of Biot Savart's law

- (i) Biot Savart's law is valid for a symmetrical current distribution.
- (ii) This law is analogous to Coulomb's law in electrostatics.
- (iii) The direction of \overrightarrow{dB} is perpendicular to both I \overrightarrow{dl} and \overrightarrow{r} .

03. Magnetic Field Due to a Straight Wire Carrying Current



$$d\mathbf{B} = \frac{\mu_0}{4\pi} \times \frac{Idl \sin \theta}{r^2}$$

$$\cos \phi = \frac{a}{r} \text{ or } r = \frac{a}{\cos \phi}$$

$$\tan \phi = \frac{1}{a}$$
 or $l = a \tan \phi$

$$dB = \frac{\mu_0}{4\pi} \frac{I(a \sec^2 \phi \, d\phi) \cos \phi}{\left(\frac{a^2}{\cos^2 \phi}\right)} = \frac{\mu_0}{4\pi} \frac{I}{a} \cos \phi \, d\phi$$

$$B = \frac{\mu_0}{4\pi} \frac{I}{a} \left(\sin \phi_1 + \sin \phi_2 \right)$$

04. Magnetic Field at a Point on the Axis of a Circular Coil Carrying Current

Plane of the coil be perpendicular to the plane of the paper and current I be flowing in there coil in the direction shown.

