

BIOLOGY

CLASS NOTES FOR CBSE

Chapter 14. Respiration in Plants

01. Introduction

All living cells require continuous supply of energy to perform various vital activities. This energy is released in controlled manner for cellular use *via* the process of respiration. The released energy is stored in cells as ATP. Thus, ATP acts as the **energy currency** of a cell.

02. Cellular Respiration

It is defined as the mechanistic breakdown of C—C bonds of complex organic compounds with the help of oxygen within a cell to release energy. These complex organic compounds are called **respiratory substrates**. In general carbohydrates, lipids and proteins can be used as respiratory substrates.

The generalised chemical reaction representing cellular respiration is as follows



03. Aerobic Cellular Respiration

This respiration is completed in two phases

(i) Glycolysis

(ii) Breakdown of pyruvic acid

Glycolysis

The word ‘Glycolysis’ is derived from two Greek words, (*Glycos*-sugar and *lysis*-splitting). Glycolysis is also called as **Embden Meyerhof-Parnas (EMP) pathway** on the name of its contributors. It is an oxidative process, in which 1 mole of glucose is converted into 2 moles of pyruvic acid. It is common to both aerobic and anaerobic respiration. Glycolysis occurs in the cytosol of all living organisms.



MISOSTUDY.COM

The Best Online Coaching for IIT-JEE | NEET Medical | CBSE INQUIRY +91 8929 803 804





Figure 12-3
Molecular Cell Biology, Sixth Edition
© 2008 W. H. Freeman and Company

Net Result of Glycolysis

The overall reaction of glycolysis is as follows



- During glycolysis, two ATP molecules are consumed in the conversion of glucose into fructose-1, 6-diphosphate. On the other hand, it produces 4 ATP and 2 NADH₂ molecules.
- The two molecules of NADH + H⁺ enter into the mitochondria and get oxidised through ETS to form 3 ATP each, i.e. a total of 6 ATP.
- Thus, the total or net gain of ATP in glycolysis process is (4 ATP produced — 2 ATP consumed) + 6 ATP from NADH₂ = **8 ATP** molecules.

NOTE **Substrate level phosphorylation** is a type of chemical reaction. In this reaction a phosphate rich compound (usually ATP) transfers one of its phosphate group to a substrate, e.g. ATP becomes ADP after transferring one of its phosphate group to the substrate. Glycolysis involves two substrate level phosphorylation reactions.


Fate of pyruvic acid

Pyruvic acid is the end product of glycolysis. Its fate depends upon the availability of oxygen in the cell. Three major ways in which cells handle pyruvic acid are lactic acid fermentation, alcoholic fermentation and aerobic respiration. The relationship between aerobic and anaerobic respiration is represented by Pfeffer-Kostychev scheme below



MISOSTUDY.COM

The Best Online Coaching for IIT-JEE | NEET Medical | CBSE INQUIRY +91 8929 803 804




Breakdown of Pyruvic Acid

The breakdown of pyruvic acid formed during glycolysis occurs in following two steps

(i) Oxidative Decarboxylation of Pyruvic Acid


The pyruvic acid formed in glycolysis moves into the mitochondria. Here, oxidative decarboxylation of pyruvic acid takes place. During this process, pyruvic acid is first decarboxylated, i.e. releases CO₂. It is then dehydrogenated by removing H-atom. Thus, 3-C pyruvic acid is converted into 2-C acetyl coenzyme-A. This process is catalysed by a large enzyme complex called **pyruvate dehydrogenase**. This enzyme complex consists of Mg²⁺, thiamine pyrophosphate, NAD, coenzyme-A and lipoic acid as cofactors. This complete process takes place in perimitochondrial, space. The reaction for this process is as follows



Acetyl Co-A formed above reacts with oxaloacetate in Krebs' cycle. Thus, this step is considered as the connective link between glycolysis and krebs' cycle.

(ii) Tricarboxylic Acid or Krebs' Cycle

This cycle is also called as **citric acid cycle**, because citric acid is the first product of this cycle. It was discovered by Sir Hans Krebs' in 1937. In eukaryotic organisms, all reactions of Krebs' cycle take place in the mitochondrial matrix. Thus, all enzymes of this cycle are also found in matrix except succinic or succinate dehydrogenase. This enzyme is found to be located in the inner membrane of mitochondria. In prokaryotes, Krebs' cycle occurs in cytoplasm. The overall reaction of aerobic degradation of pyruvic acid is as follows. (This includes oxidative decarboxylation and TCA)



MISOSTUDY.COM

The Best Online Coaching for IIT-JEE | NEET Medical | CBSE INQUIRY +91 8929 803 804