

Complete
CHEMISTRY

IIT-JEE · NEET · CBSE eBOOKS

CLASS 11 & 12th



Learning Inquiry
8929 803 804

CLASS 11th


Hydrocarbons

misostudy



01. Introduction

Organic compounds composed of only carbon and hydrogen are called hydrocarbons.






02. Saturated Hydrocarbons

These constitute a homologous series having general formula C_nH_{2n+2} (n may have value 1,2,3,4).


The saturated hydrocarbons are called **paraffins** (Latin: parum= little; affinity= affinity) as they are relatively inert toward chemical reagents. In IUPAC nomenclature, paraffins are termed alkanes. Alkanes have following structural characteristics,

- Every carbon atoms is Sp^3 hybridized, its four bonding orbitals are directed toward the four corner of a regular tetrahedron.
- All the carbon-carbon and carbon-hydrogen bonds are strong sigma. The carbon-carbon bond is formed from the overlap of Sp^3 orbitals, one from each carbon atom. All carbon-hydrogen bonds result in overlap of Sp^3 hybrid orbitals, one from each carbon atom. All carbon-hydrogen bonds result in overlap of Sp^3 hybrid orbital from carbon and s-orbital from hydrogen.
- The bond lengths between carbon-hydrogen are 1.54Å and 1.112Å respectively.
- The bond angles in alkanes are tetrahedral angles having a value of 109.5° ($109^\circ28'$).






The structure of methane molecule can be represented by (a), (b) and (c).
Fig. (a) is the **tetrahedral structure** of methane in which.



Structure Isomerism in alkanes











Alkanes exhibit chain isomerism. The first three members, *viz.*, methane, ethane and propane do not exhibit isomerism as they can be represented by only one structural formula. Butane has two chain isomers.




With increase in the number of carbon atoms in the molecule, the number of chain isomers also increases.

Alkane	C_3H_{12}	C_6H_{14}	C_7H_{16}	C_8H_{18}	$\text{C}_{10}\text{H}_{22}$
No. of possible isomers	3	5	9	18	75

Greater the branching greater the stability; so increasing order of stability is:
n-pentane < iso-pentane < neo-pentane;

Preparation of Alkanes**(i) From unsaturated hydrocarbons****(ii) From alkyl halides****(a) Reduction :****(b) Wurtz reaction :****(iii) From carboxylic acids****(a) By decarboxylation of carboxylic acids :****(b) Kolbe's electrolytic method :**

Mechanism : The reaction follows a free-radical mechanism :



At anode :

