

Complete
CHEMISTRY

IIT-JEE · NEET · CBSE eBOOKS

CLASS 11 & 12th



Learning Inquiry
8929 803 804

CLASS 11th

P-Block Elements

misostudy



01. Physical Properties

(i) Atomic Size

Atomic Number	Atomic Radil	Element	Metalic Character
5	85 pm	B	Non-metal
13	143 pm	Al	Metal
31	135 pm	Ga	Metal
49	167 pm	In	Metal
81	170 pm	Tl	Metal

(ii) Electronegativity

Atoms	B	Al	Ga	In	Tl
Electronegativity	2	1.5	1.6	1.7	1.8

Due to poor shielding of 3d and 4d and lanthanoid contraction.

(iii) Ionization Energy

$B > Tl > Ga > Al > In$

(iv) Oxidation States

Stability of +3 oxidation state decreases down the group due to inert pair effect, as well stability of +1 oxidation state increases.

(v) Melting Point

$B > Al > Tl > In > Ga$

Ga has on unused structure. It congests if only Ga_2 molecules it has, thus low melting point.


(vi) Boiling Point

$B > Al > Ga > In > Tl$

NOTE ☞ Indium in +1 oxidation state is reducing agent.

(i) Reactivity towards air




B_2O_3 — Acidic Oxide
 AlO_3 — Amphoteric Oxide
 Ga_2O_3 — Amphoteric Oxide
 In_2O_3 — Basic Oxide
 Tl_2O_3 — Basic Oxide



B_2O_3 is called boric anhydride as it is anhydride of boric acid

(ii) **Reactivity towards acids**


Boron is not affected by acids agents like HCl and dil. H_2SO_4 while all other elements react with conc. H_2SO_4 and HNO_3 . Ga and Al develop protective layer of oxide with conc. HNO_3 .

(iii) **Reactivity towards alkalis**

Except indium and thallium all other elements react with alkali solutions




Example, Al also reacts with aq. Alkali and liberates dihydrogen.


(iv) **Reactivity towards halogens**

Trihalides are formed when these elements react with halogens. All these halides exist as discrete molecular species which are sp^2 hybridised and covalently bonded. TlI_3 is unstable.


AlCl_3 achieves stability by forming a dimer.



Acidic strength is inversely proportional to back-bonding, as back-bonding decreases from BF_3 to BI_3 as given below



Hence, Lewis acidic strength will increase as under



$p\pi$ - $p\pi$ back-bonding is strongest in BF_3 because both B and F involve $2p$ orbital in back-bonding. Stability of halides in +3 oxidation state decreases down the group.

Anomalous Behaviour of Boron

- (i) Boron has very small atomic radii, hence greater nuclear attraction on the outermost electrons. It has very high ionisation energy. This gives boron distinctly non-metallic character while the rest are metals.
- (ii) Boron has maximum covalency of four due to non-availability of d -orbitals while the rest have maximum covalency of six.
- (iii) Boron alone exhibits allotropy.
- (iv) Boron shows +3 oxidation state while others can show +1 and +3 oxidation states.

Comparison of Boron and Aluminium

- (i) Same Electronic Configuration
- (ii) Same Valency
- (iii) Same Oxidation State