



IIT-JEE · CBSE **eBOOKS**

CLASS 11 & 12th



Learning Inquiry  
8929 803 804

CLASS 12th

# Determinants

misostudy



## 01. Determinants

### DEFINITION

Every square matrix can be associated to an expression or a number which is known as its determinant. If  $A = [a_{ij}]$  is a square matrix of order  $n$ , then the determinant of  $A$  is denoted by  $\det A$  or,  $|A|$  or,

$$\begin{vmatrix} a_{11} & a_{12} & \dots & a_{1j} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2j} & \dots & a_{2n} \\ \vdots & \vdots & & & & \\ a_{i1} & a_{i2} & \dots & a_{ij} & \dots & a_{in} \\ \vdots & \vdots & & \vdots & & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nj} & \dots & a_{nn} \end{vmatrix}$$

### DETERMINANT OF A SQUARE MATRIX OF ORDER 1

If  $A = [a_{11}]$  is a square matrix of order 1, then the determinant of  $A$  is defined as

$$|A| = a_{11} \quad \text{or,} \quad |a_{11}| = a_{11}$$

### DETERMINANT OF A SQUARE MATRIX OF ORDER 2

The determinant of a square matrix of order 2 is equal to the product of the diagonal elements minus the product of off-diagonal elements.

### DETERMINANT OF A SQUARE MATRIX OF ORDER 3

If  $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$  is a square matrix of order 3, then the expression

$$a_{11} a_{22} a_{33} + a_{12} a_{23} a_{31} + a_{13} a_{32} a_{21} - a_{11} a_{23} a_{32} - a_{22} a_{13} a_{31} - a_{12} a_{21} a_{33}$$

is defined as the determinant of  $A$  i.e.

$$\begin{aligned} |A| &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ &= a_{11} a_{22} a_{33} + a_{12} a_{23} a_{31} + a_{13} a_{32} a_{21} - a_{11} a_{23} a_{32} - a_{22} a_{31} a_{13} - a_{33} a_{12} a_{21} \dots \text{(ii)} \end{aligned}$$

$$\begin{aligned} \text{or,} \quad |A| &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ \Rightarrow |A| &= a_{11} (a_{22} a_{33} - a_{23} a_{32}) - a_{12} (a_{33} a_{21} - a_{23} a_{31}) + a_{13} (a_{32} a_{21} - a_{22} a_{31}) \\ \Rightarrow |A| &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \quad [\text{Using notation given in (i)}] \\ \Rightarrow |A| &= (-1)^{1+1} a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} + (-1)^{1+2} a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + (-1)^{1+3} a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \end{aligned}$$

Thus the determinant of a square matrix of order 3 is the sum of the product of elements  $a_{1j}$  in first row with  $(-1)^{1+j}$  times the determinant of a  $2 \times 2$  sub-matrix obtained by leaving the first row and column passing through the element.

- NOTE** ✍
1. Only square matrices have determinants. The matrices which are not square do not have determinants.
  2. The determinant of a square matrix of order 3 can be expanded along any row or column.
  3. If a row or a column of a determinant consists of all zeros, then the value of the determinant is zero.

## DETERMINANT OF A SQUARE MATRIX OF ORDER 4 OR MORE

To evaluate the determinant of a square matrix of order 4 or more we follow the same procedure as discussed in evaluating the determinant of a square matrix of order 3.

## 02. Minors and Cofactors

### Minor

Let  $A = [a_{ij}]$  be a square matrix of order  $n$ . Then the minor  $M_{ij}$  of  $a_{ij}$  in  $A$  is the determinant of the square sub-matrix of order  $(n-1)$  obtained by leaving  $i^{th}$  row and  $j^{th}$  column of  $A$ .

### Cofactor

Let  $A = [a_{ij}]$  be a square matrix of order  $n$ . Then, the cofactor  $C_{ij}$  of  $a_{ij}$  in  $A$  is equal to  $(-1)^{i+j}$  times the determinant of the sub-matrix of order  $(n-1)$  obtained by leaving  $i^{th}$  row and  $j^{th}$  column of  $A$ .

It follows from this definition that

$$C_{ij} = \text{Cofactor of } a_{ij} \text{ in } A = (-1)^{i+j} M_{ij}, \text{ where } M_{ij} \text{ is minor of } a_{ij} \text{ in } A.$$

Thus, we have

$$C_{ij} = \begin{cases} M_{ij} & \text{if } i+j \text{ is even} \\ -M_{ij} & \text{if } i+j \text{ is odd} \end{cases}$$

## EXPANSION OF DETERMINANT USING MINORS/CO-FACTORS

The value of determinant is defined as the sum of the product of elements of any row (column) by their corresponding co-factors.

$$\begin{aligned} \text{Let } \Delta &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \\ &= \underbrace{a_{11} C_{11} + a_{12} C_{12} + a_{13} C_{13}}_{\text{expanding along } R_1} \\ &= a_{11} a_{22} a_{33} - a_{11} a_{32} a_{23} + a_{12} a_{31} a_{23} - a_{12} a_{21} a_{33} + a_{13} a_{21} a_{32} - a_{13} a_{31} a_{22} \end{aligned}$$

Therefore the value of the determinant can be obtained

as  $\Delta = a_{11} C_{11} + a_{12} C_{12} + a_{13} C_{13}$  (expanding along  $R_1$ )

or  $\Delta = a_{12} C_{12} + a_{22} C_{22} + a_{32} C_{32}$  (expanding along  $C_2$ )

or  $\Delta = a_{31} C_{31} + a_{32} C_{32} + a_{33} C_{33}$  (expanding along  $R_3$ )

In general, expanding along  $i^{th}$  row we get

$$\Delta = a_{i1} C_{i1} + a_{i2} C_{i2} + \dots + a_{in} C_{in} = \sum_{k=1}^n a_{ik} C_{ik} \quad (\text{for all } i = 1, 2, \dots, n)$$

And expanding along  $j^{th}$  column, we get

or  $\Delta = a_{1j} C_{1j} + a_{2j} C_{2j} + \dots + a_{nj} C_{nj} = \sum_{k=1}^n a_{kj} C_{kj} \quad (\text{for all } j = 1, 2, \dots, \text{or } n)$

- NOTE** 1. The expansion generates same value irrespective of its performance through any row or column.  
 2. The expansion contains  $3!$  i.e., 6 terms which is the number of permutations of 1, 2, 3 in a line.

$$\text{i.e., } \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = \underbrace{(a_1 b_2 c_3 + b_1 c_2 a_3 + c_1 a_2 b_3)}_{\text{positive zone}} - \underbrace{(a_1 b_3 c_2 + b_1 a_2 c_3 + c_1 a_3 b_2)}_{\text{-ative zone}}$$

3. Each term is product of three entries of the determinant.
4. 3 terms are positive, 3 other are negative (even and odd permutations).
5. Each entry of the determinant  $\Delta$  once appears in the positive zone and once in the negative zone. For instance  $a_1$  appears as in the first and fourth term.
6. A square matrix is a singular matrix if its determinant is zero. Otherwise, it is a non-singular matrix.

### 03. Properties of Determinants

**Property 1**

Let  $A = [a_{ij}]$  be a square matrix of order  $n$ , then the sum of the product of elements of any row (column) with their cofactors is always equal to  $|A|$  or,  $\det(A)$  i.e.

$$\sum_{j=1}^n a_{ij} C_{ij} = |A| \quad \text{and} \quad \sum_{i=1}^n a_{ij} C_{ij} = |A|.$$

**Property 2**

Let  $A = [a_{ij}]$  be a square matrix of order  $n$ , then the sum of the product of elements of any row (column) with the cofactors of the corresponding elements of some other row (column) is zero i.e.

$$\sum_{j=1}^n a_{ij} C_{ij} = 0 \quad \text{and} \quad \sum_{i=1}^n a_{ij} C_{ik} = 0.$$

**Property 3**

Let  $A = [a_{ij}]$  be a square matrix of order  $n$ , then  $|A| = |A^T|$ .

or, the value of a determinant remains unchanged if its rows and columns are interchanged.