

MATHEMATICS

Live eBook



01. Joint Equation of A Pair of Straight Lines

The joint equation of the straight lines $a_1 x + b_1 y = 0$ and $a_2 x + b_2 y = 0$ is of the form $ax^2 + 2hxy + by^2 = 0$ which is known as the homogeneous equation of second degree. Thus, the combined equation of a pair of straight lines passing through the origin is a homogeneous equation of second degree.

Consider the equation $ax^2 + 2hxy + by^2 = 0$. This equation can be re-written as

$$b\left(\frac{y}{x}\right)^2 + 2h\left(\frac{y}{x}\right) + a = 0 \quad [\text{Dividing throughout by } x^2]$$
$$\Rightarrow \frac{y}{x} = \frac{-h \pm \sqrt{h^2 - ab}}{b}$$
$$\Rightarrow y = \left(\frac{-h \pm \sqrt{h^2 - ab}}{b}\right)x$$
$$\Rightarrow y = m_1 x, \quad y = m_2 x, \quad \text{where}$$
$$m_1 = \frac{-h + \sqrt{h^2 - ab}}{b} \quad \text{and} \quad m_2 = \frac{-h - \sqrt{h^2 - ab}}{b}$$

The value of m_1 and m_2 are (i) real and distinct if $h^2 > ab$ (ii) real and equal if $h^2 = ab$, and (iii) imaginary if $h^2 < ab$.

Also, $m_1 + m_2 = -\frac{2h}{b}$ and, $m_1 m_2 = \frac{a}{b}$

Thus, $ax^2 + 2hxy + by^2 = 0$ represents a pair of straight lines $y = m_1 x$ and $y = m_2 x$ which are real and distinct if $h^2 > ab$. Also,

$$m_1 + m_2 = -\frac{2h}{b} = -\frac{\text{Coefficient of } xy}{\text{Coefficient of } y^2}$$

and, $m_1 m_2 = \frac{a}{b} = -\frac{\text{Coefficient of } x^2}{\text{Coefficient of } y^2}$

02. Angle Between The Pair of Lines

The acute angle between the two lines represented by $ax^2 + 2hxy + by^2 = 0$ & are given as $y = m_1 x$ (slope = m_1) & $y = m_2 x$ (slope = m_2) is

$$\tan \theta = \left| \frac{m_2 - m_1}{1 + m_1 m_2} \right|$$
$$\tan \theta = \frac{|m_2 - m_1|}{|1 + m_1 m_2|} = \frac{\sqrt{(m_2 - m_1)^2}}{|1 + m_1 m_2|} = \frac{\sqrt{m_2^2 + m_1^2 - 2m_1 m_2}}{|1 + m_1 m_2|}$$
$$= \frac{\sqrt{m_1^2 + m_2^2 + 2m_1 m_2 - 4m_1 m_2}}{|1 + m_1 m_2|}$$
$$= \frac{\sqrt{(m_1 + m_2)^2 - 4m_1 m_2}}{|1 + m_1 m_2|}$$

$$\begin{aligned}
&= \frac{\sqrt{\left(-\frac{2h}{b}\right)^2 - 4\frac{a}{b}}}{\left|1 + \frac{a}{b}\right|} = \frac{\sqrt{\frac{4h^2}{b^2} - \frac{4a}{b}}}{\frac{|a+b|}{|b|}} \\
&= \frac{\frac{\sqrt{4h^2 - 4ab}}{|b|}}{\frac{|a+b|}{|b|}} = \frac{2\sqrt{h^2 - ab}}{|a+b|} = \frac{2\sqrt{h^2 - ab}}{|a+b|}
\end{aligned}$$

Lines are parallel iff they are coincident
iff $h^2 = ab$
($\tan \theta = 0 \Rightarrow \theta = 0$)

Lines are perpendicular iff $m_1 m_2 = -1$
iff $\frac{a}{b} = -1$
or $a = -b$ or $a + b = 0$
 $a + b = 0$

